

Baseline urinalysis results in 32 healthy Antillean manatees (*Trichechus manatus manatus*)

Lesly J. Cabrias-Contreras MVZ, Dipl.

Roberto Sánchez-Okrucky MVZ

Dalila Caicedo-Herrera Bióloga Marina

Laura Jaramillo-Ortiz MV, MSc

Francisco de la Rosa DMV

Ana C. Negrete-Philippe MVZ

Danilo Cruz-Martínez MVZ, PhD

Antonio L. Rivera-Guzmán DMV

Antonio Mignucci-Giannoni VT, PhD

From the Caribbean Manatee Conservation Center, Inter-American University of Puerto Rico, San Juan, Puerto Rico (Cabrias-Contreras, Rivera-Guzmán, Mignucci-Giannoni); Dolphin Discovery Group, Cancún, Quintana Roo, México (Sánchez-Okrucky, Cruz-Martínez); Fundación Omacha, Bogotá, Colombia (Caicedo-Herrera, Jaramillo-Ortiz); Acuario Nacional de la República Dominicana, Santo Domingo, República Dominicana (de la Rosa); Grupo Xcaret, Playa del Carmen, Quintana Roo, México (Negrete-Philippe); Center for Conservation Medicine and Ecosystem Health, Ross University School of Veterinary Medicine, Basseterre, St. Kitts (Mignucci-Giannoni).

Address correspondence to Dr. Mignucci-Giannoni (mignucci@manatipr.org).

OBJECTIVE

To describe results of free-catch urine samples collected from Antillean manatees (*Trichechus manatus manatus*) under human care in the Caribbean.

ANIMALS

32 Antillean manatees in 5 Caribbean oceanaria and rescue centers.

PROCEDURES

Urine samples were obtained by opportunistic free catch during physical examination or through the use of operant conditioning procedures. Urinalyses consisted of macro- and microscopic evaluations, biochemical analyses with test strips, and refractometry. Results were compared for manatees grouped on the basis of age, sex, and habitat.

RESULTS

Urine samples were typically clear, straw colored, and alkaline (mean pH, 8.0); had a urinous odor and low specific gravity (mean, 1.010); and had results on qualitative test strips that were consistently negative for the presence of glucose, bilirubin, ketones, proteins, nitrites, RBCs, and WBCs. Microscopically, the mean \pm SD number of RBCs and WBCs/hpf was 0.5 ± 0.3 RBCs/hpf and 1.1 ± 1.5 WBCs/hpf. Findings of some epithelial cells and crystals were typical. Spermatozoa were found in urine from 1 of 15 sexually mature males, and parasite larvae and eggs were found in urine from 2 manatees.

CONCLUSIONS AND CLINICAL RELEVANCE

Results of the present study yielded the first compilation of baseline urinalysis values in healthy Antillean manatees under human care, which, when combined with physical examination and other diagnostic procedures, can help in monitoring the health of these animals. We encourage the use of free-catch urine collection methods, as used in the present study, for routine urinalyses of manatees under human care in zoos, aquaria, or rescue centers. (*J Am Vet Med Assoc* 2021;258:416–424)

Antillean manatees (*Trichechus manatus manatus*) are endangered,^a euryhaline, herbivorous aquatic mammals that live in marine, estuarine, and freshwater ecosystems of the Caribbean.¹ Manatees have been kept under human care since the 1950s,² and although considerable attention has been given to their care (including veterinary diagnostic procedures and treatments) during the past 30 years,^{3–7} urinalysis is infrequently performed in manatees housed at aquariums and zoos.⁸ In previous studies,^{8–10,b} urine samples from Florida manatees (*Trichechus manatus latirostris*) and Amazonian manatees (*Trichechus inunguis*) were examined. However, Antillean manatees and Florida manatees are subspecies that differ morphologically, physiologically, and genetically from each other and from Amazonian manatees,^{11–14} and there is no published information on results of urinalysis for Antillean manatees. Our aim of the study reported here was to describe the macroscopic, microscopic, and biochemical characteristics of

free-catch urine samples collected through the use of noninvasive medical operant conditioning in Antillean manatees under human care in the Caribbean.

Materials and Methods

Animals and facilities

Thirty-two Antillean manatees under human care at 5 oceanaria and rescue center facilities in Colombia, Dominican Republic, Mexico, and Puerto Rico were enrolled in the study, which was approved by the Inter-American University Institutional Animal Care and Use Committee and conducted in accordance with the US Animal Welfare Act. On the basis of results from physical and hematologic examinations, the manatees were determined to have been clinically normal at the time of urine sampling between January 16, 2016, and November 20, 2019. The manatees were grouped on the basis of sex, age (ie, calf, juvenile, subadult, or adult¹⁵), and housing environment (ie, freshwater, estuarine, or saltwater). Water salinity at each facility was measured with a salinity refractometer.^c All facilities were located in the tropical zone, in Caribbean waters with air tempera-

ABBREVIATIONS

ppt Parts per thousand
SG Specific gravity

tures between 25°C and 29°C (77°F to 84.2°F) and water temperatures between 26°C and 29°C (78.8°F to 84.2°F). Diet consisted of aquatic plants and vegetables for adult, subadult, and juvenile manatees. For calves, natural or artificial milk formula with small inclusions of vegetables was fed, and depending on each calf's health and age, this diet was supplemented with various multivitamins and with biscuits formulated for folivore primates.^d

Sample collection

Before a manatee's first feeding of the day, a mid-stream urine sample was obtained by opportunistic free catch during a veterinary examination or trained medical behaviors^{16–18,e} (**Figure 1**). Urine collection from manatees in Puerto Rico, Colombia, and the Dominican Republic was strictly opportunistic during veterinary evaluations in an empty tank devoid of water or on land along the side of a pond enclosure. Manatees were subject to regular physical examinations on the bottom of the empty tank or on land, during which the animal would typically lie in dorsal recumbency, facilitating urine collection. In Mexico, manatees were trained to urinate on command through operant conditioning, similar to conditioning used in dolphins,¹⁹ except that instead of a hand signal for the urination command, light hand pressure was applied in a sustained manner over the area of the urinary bladder, just cranial to the anus. This pressure would trigger sufficient urination. Because sampling took several minutes in some instances, the trained behavior for urine sample collection included allowing the manatee to breathe without compro-

mising the cleanliness of the genital area, which had been cleaned with isopropyl alcohol and allowed to air-dry before each sample collection.

Urine samples were collected in sterile, 120-mL polypropylene containers^f or with a sterile 10- to 20-mL plastic syringe. Following collection, the container or syringe was labeled with the manatee's name and the date and time of collection. Urine samples were transported to the laboratory on ice and stored, when needed, at 4°C to 5°C (39.2°F to 41°F) until urinalysis was completed within 1 hour of collection.

Urinalysis

Urinalysis consisted of gross examination followed by biochemical analyses and microscopic evaluation of urine sediment as described previously.²⁰ A standardized format for marine mammals^g was used to record data, which were then transferred to a spreadsheet^h at the end of each sampling and analysis day. Prior to analysis of results, a comparison was made between the recorded information on the urinalysis form versus the computerized spreadsheet to identify and correct any transcription errors, if present.

Gross examination—A sterile, polymethylpentene plastic 50-mL graduated cylinderⁱ was used to measure the volume of each urine sample. Physical characteristics of each sample were described, including color (ie, colorless, pale yellow, yellow, dark yellow, dark brown-yellow, pale green-yellow, green-yellow, dark green-yellow, straw, ochre, red-brown, or red), appearance (ie, clear, hazy, or slightly cloudy), and odor (ie none, urinoid [strong urine smell], fruity or



Figure 1—Representative photograph of the use of operant conditioning in collecting urine from 1 of the 12 Antillean manatees (*Trichechus manatus manatus*) under human care in Mexico between January 19 and July 20, 2017. The manatee in the image is an adult female.

sweet, fruity fishy [mix of sweet and fish smell], fishy, grassy [smell of grass], pungent [having a sharply strong odor], sulfuric [rotten egg smell], aromatic [fragrant smell], fecal [smelling of feces], or ammoniacal [long-standing ammonia smell]). Because physical characteristics of urine are subjective, the evaluators were trained in a veterinary laboratory and practiced gross urine examination on urine collection from domestic mammals and dolphins.

Biochemical examination—Following gross examination, each urine sample was transferred to a sterile, 15-mL conical polypropylene centrifuge tube^j for biochemical analyses with 3 different types of test strips. One type of test strip^k was used to assess urine SG, pH, and concentrations of glucose, bilirubin, urobilinogen, ketones, nitrite, protein, RBCs, and WBCs. A second type of test strip^l was also used to assess urine pH, and a third type of test strip^m was used to assess urine concentrations of creatinine and microalbumin. Each test strip was immersed in the urine sample, then removed immediately. Excess urine was removed from the strip by touching a paper towel to the edge of the strip. The color change for each urinalysis parameter was compared with a colorimetric scale supplied with the respective test strip kit, and the results were recorded. Qualitative results for whether analytes were present versus absent (negative) were recorded, and if present, semiquantification of results to the extent feasible with the test strips was recorded (eg, urine glucose concentration: negative vs 100, 250, 500, 1,000, or > 2,000 mg/dL). Detection of semiquantitative trace concentration urobilinogen (ie, ≤ 0.2 mg/dL) was considered a negative result for the detection of an abnormal concentration of urobilinogen. Because a study¹⁹ of dolphins shows that measurements of urine SG were inaccurate when determined with the same type of test strip^k used in the present study, we also measured urine SG using a veterinary clinical refractometerⁿ that was calibrated with distilled water.

Microscopic examination—To extract urinary sediment from each sample, approximately 3 mL of urine was retained in the conical tube and centrifuged^o for 2.5 minutes at 1,163 \times g. Pipette aspiration was used to discard most of the supernatant, leaving a volume of approximately 0.5 to 1 mL of supernatant in which the sediment pellet was resuspended. A drop of the resuspended sediment was placed on a glass slide, covered with a cover slip, and examined with light microscopy at an absolute magnification of 100X and 400X (hpf) to assess for WBCs, RBCs, epithelial cells (transitional, squamous, or tubular), casts (hyaline, granular, waxy, epithelial, or fatty), crystals (calcium oxalate, uric acid, triple phosphate, calcium phosphate, amorphous phosphate, or urate), bacteria, yeast or fungi, spermatozoa, and parasites or parasite eggs. For each urine sample, the mean numbers of microscopy elements of interest (eg, WBCs, RBCs, and epithelial cells)/10 hpf (400X) were determined, recorded, and then qualitatively categorized as none (mean, 0 elements/hpf), rare (mean,

> 0 to ≤ 2.5 elements/hpf), few (mean, > 2.5 to < 5.0 elements/hpf), some (mean, ≥ 5.0 to < 7.5 elements/hpf), or many (mean, ≥ 7.5 elements/hpf). Qualitative results for mucus were defined as none (not observed), rare (occasionally present), few (mildly present), some (moderately present), or many (abundantly present).

Statistical analysis

Descriptive statistics were calculated and analyzed with statistical software.^p Qualitative results were reported as numbers and percentages, and quantitative results were reported as range and mean \pm SD.²¹⁻²⁴ Results were assessed for normality of distribution with the Shapiro-Wilk test, and paired 2-tailed *t* tests were used to evaluate for differences in mean results for manatees grouped on the basis of age, sex, and habitat. Values of $P \leq 0.05$ were considered significant.

Results

Animals

There were 32 manatees (21 males [15 sexually mature] and 11 females [8 sexually mature]) under human care at 5 oceanaria and rescue center facilities in Colombia ($n = 9$), Dominican Republic (2), Mexico (12), and Puerto Rico (9). Of the 32 manatees, 6 were calves (age, ≤ 1 year), 3 were juveniles (age, 2 to 4 years), 3 were subadults (age, 5 to 9 years), and 20 were adults (age, ≥ 10). Of the 32 manatees, 13 were in open marine systems (salinity range, 31 to 35 ppt), 6 were in open cenote estuarine systems (salinity range, 17 to 22 ppt), and 13 were in freshwater cement tanks or river systems (salinity, 0 ppt).

Urine samples

Free-catch urine samples were obtained by opportunistic free catch during veterinary examinations for 20 manatees and through operant conditioning procedures for 12 manatees. For the manatees with which operant conditioning was used, urination occurred within 5 to 8 minutes after the start of the requesting procedure. Overall, the mean \pm SD urine sample volume was 5.4 ± 1.53 mL (range, 3 to 9 mL), and the mean \pm SD duration from collection to urinalysis was 19.5 ± 6.8 minutes (range, 8 to 35 minutes).

Gross examination—The color of most urine samples was straw (22/32 [69%]), followed by pale yellow (6/32 [19%]) or colorless (4/32 [13%]; **Table 1**). In addition, the appearance of most samples was clear (21/32 [66%]), and the most common odor of samples was urinoid (strong urine smell; 25/32 [78%]).

Biochemical examination—Qualitative results of urinalyses were predominantly negative for biochemical analytes assessed with test strips; however, semiquantitative trace concentrations were detected for urobilinogen (0.2 mg/dL; $n = 32/32$), creatinine (0.1 mg/dL; 17/17), and microalbumin (10 mg/dL; 17/17; **Table 2**). The microalbumin-to-creatinine ratio was 100 mg/g for all 17 manatees from which urine samples were evaluated for both analytes.

Overall, the mean \pm SD SG measured by refractometry was 1.010 ± 0.007 and did not differ substantially between males (1.011 ± 0.008) and females (1.009 ± 0.005). When considered by manatee age group, the mean \pm SD urine SG was 1.009 ± 0.004 for calves ($n = 6$), 1.010 ± 0.011 for juveniles (3), 1.005 ± 0.001 for subadults (3), and 1.011 ± 0.008 for adults (20). On the basis

of manatee habitat, the mean \pm SD urine SG was 1.012 ± 0.009 for those in marine habitats ($n = 13$), 1.011 ± 0.005 for those in estuarine habitats (6), and 1.008 ± 0.006 for those in freshwater habitats (13). The small number of manatees in each of the various categories precluded further statistical analyses regarding urine SG.

Overall, the mean \pm SD urine pH was 8.0 ± 0.9 and did not differ substantially between males (7.9 ± 1.1) and females (8.0 ± 0.7). By age group, the mean \pm SD urine pH was 7.8 ± 0.7 for calves, 6.7 ± 1.5 for juveniles, 8.0 ± 0.0 for subadults, and 8.2 ± 0.9 for adults. On the basis of manatee habitat, the mean \pm SD urine pH was 8.3 ± 0.9 , 8.0 ± 0.8 , or 7.6 ± 1.0 for those in marine, estuarine, or freshwater habitats, respectively. Similar to findings for SG, the small number of manatees in each category precluded further statistical analyses regarding urine pH.

Microscopic examination—In urinary sediments of the 32 manatees, the mean \pm SD number of RBCs/hpf was 0.5 ± 0.3 (range, 0 to 3 RBCs/hpf), and the mean number of WBCs/hpf was 1.1 ± 1.5 (range, 0 to 8 WBCs/hpf), respectively. Qualitatively, epithelial cells were observed in urinary samples from 28 manatees, with transitional cells less commonly present (1/32 [3%]), compared with squamous cells (21/32 [66%]), which were more commonly identified in samples from females (11/11 [100%]) versus males (10/21 [48%]; **Table 3**). Urinary crystals were observed in samples from 13 manatees, with calcium oxalate monohydrate crystals most commonly identified among the affected samples ($n = 4$). There were rare observations of mucus, erythrocyte casts, hyphae, sperm, or parasite larvae and eggs, and these were found only in samples from adult male manatees. For instance, casts were present in urine samples from only 2 male manatees housed in an estuarine system in Mexico, and both manatees had RBC-type casts. In addition, these 2 males had unidentified nematode-type larvae and eggs observed in their urine. A summary of key urinalysis find-

Table 1—Gross characteristics of free-catch urine samples obtained from 32 Antillean manatees (*Trichechus manatus manatus*) under human care in 5 Caribbean oceanaria and rescue centers between January 16, 2016, and November 20, 2019.

Characteristic	No. (%) of urine samples
Color	
Straw	22 (69)
Pale yellow	6 (19)
Colorless	4 (13)
Yellow	0
Dark yellow	0
Dark brown-yellow	0
Pale green-yellow	0
Green-yellow	0
Dark green-yellow	0
Ochre	0
Red-brown	0
Red	0
Appearance	
Clear	21 (66)
Slightly cloudy	9 (28)
Hazy	2 (6)
Odor	
Urinoïd	25 (78)
Grassy	4 (13)
Fruity sweet	2 (6)
Sulfuric	1 (3)
None	0
Fruity fishy	0
Fishy	0
Pungent	0
Aromatic	0
Fecal	0
Ammoniacal	0

Table 2—Summary urinalysis results obtained with urine test strips and refractometry used to evaluate the urine samples from the 32 Antillean manatees described in Table 1.

Parameter	No. of samples				
	Tested	Negative	Positive	Range	Mean \pm SD
Glucose (mg/dL)*	32	32	0	—	—
Bilirubin (mg/dL)*	32	32	0	—	—
Urobilinogen (mg/dL)*	32	32	0	0.2†	0.2 \pm 0†
Ketone (mg/dL)*	32	32	0	—	—
Nitrite (mg/dL)*	32	32	0	—	—
Protein (mg/dL)*	32	32	0	—	—
RBCs (cells/ μ L)*	32	32	0	—	—
WBCs (cells/ μ L)*	32	32	0	—	—
SG‡	32	—	—	1.002–1.030	1.010 \pm 0.007
pH§	32	—	—	5.0–9.5	8.0 \pm 0.9
Microalbumin (mg/L)	17	0	17	10	10.0 \pm 0
Creatinine (g/L)	17	0	17	0.1	0.1 \pm 0
Microalbumin-to-creatinine ratio (mg/g)	17	—	—	100	100 \pm 0

*Measured with urine test strips.^k †The finding was considered a negative result for the detection of abnormal urobilinogen concentration in urine. ‡Measured by refractometry. §Measured with urine test strips. || Measured with urine test strips.^m
— = Not applicable.

Table 3—Qualitative findings from microscopic examination of the sediments in urine samples from the 32 manatees described in Table 1.

Sediment elements of interest	No. of samples	No. (%) of samples per qualitative category*				
		None	Rare	Few	Some	Many
Epithelial cells	32	4 (13)	6 (19)	9 (28)	12 (38)	1 (3.1)
Casts	32	30 (94)	2 (6)	0	0	0
Crystals	32	19 (60)	4 (13)	0	9 (28)	0
Bacteria	32	12 (38)	3 (10)	13 (41)	2 (6)	2 (6)
Yeast or fungi	32	30 (94)	0	2 (6)	0	0
Spermatozoa	15	14 (93)	1 (7)	0	0	0
Mucus	32	31 (97)	1 (3)	0	0	0
Parasite or ova	32	30 (94)	2 (6)	0	0	0

*For each urine sample, the mean numbers of each of the urine sediment elements of interest (eg, WBCs, RBCs, and epithelial cells)/10 hpf [400X] were determined, recorded, and then qualitatively categorized as none (mean, 0 elements/hpf), rare (mean, > 0 to ≤ 2.5 elements/hpf), few (mean, > 2.5 to < 5.0 elements/hpf), some (mean, ≥ 5.0 to < 7.5 elements/hpf), or many (mean, ≥ 7.5 elements/hpf).

Table 4—Summary of key urinalysis parameters with baseline results considered clinically normal in the 32 Antillean manatees described in Table 1.

Parameter	Baseline result
Gross examination	
Color	Straw to colorless
Appearance	Clear
Odor	Urinoid to grassy
SG	1.003–1.017
pH	7.1–8.9
Urine test strip assessments	
Glucose (mg/dL)	Negative
Bilirubin (mg/dL)	Negative
Urobilinogen (mg/dL)	≤ 0.2
Ketone (mg/dL)	Negative
Nitrite (mg/dL)	Negative
Protein (mg/dL)	Negative
RBCs (cells/μL)	Negative
WBCs (cells/μL)	Negative
Microalbumin (mg/L)	≤ 10
Creatinine (g/L)	≤ 0.1
Microalbumin-to-creatinine ratio (mg/g)	100
Microscopy of urine sediment	
WBCs (cells/hpf [400X])	0–2
RBCs (cells/hpf [400X])	0–1
Epithelial cells*	None to some
Casts*	None
Crystals*	None to some
Bacteria*	None to few
Yeast or fungi*	None
Spermatozoa*	None
Mucus†	None
Parasites or ova of parasites*	None

*Reported as the qualitative categories described in Table 3. †Qualitative results for mucus were defined as none (not observed), rare (occasionally present), few (mildly present), some (moderately present), many (abundantly present).

ings for gross, biochemical, and microscopic characteristics identified in the 32 healthy Antillean manatees under human care was compiled (**Table 4**).

Discussion

Urinalysis is a basic veterinary diagnostic tool used to evaluate renal function.²⁵ It is easy, quick, and inexpensive, and we used noninvasive techniques to obtain urine

samples from the manatees in the present study. Together with physical examinations, hematologic and serum biochemical analyses, and coprologic tests, urinalysis is a key diagnostic tool for evaluating patients, including routinely evaluating wild animals maintained under human care.²⁶ Many zoos and aquaria have increasingly relied on training animal behaviors conducive to medical procedures through operant conditioning. Because cystocentesis or urinary catheterization would require patient restraint and could result in patient resistance, injury to the staff and patient, and less cooperation from the patient in future events, the use of operant conditioning to obtain diagnostic samples is less invasive, minimizes stress during sampling, and increases the ease in which basic diagnostic procedures can be performed,¹⁸ including in manatees.^{16,17} If renal dysfunction is suspected on the basis of results from samples collected with operant conditioning, further tests (eg, assessments of electrolytes and urea waste product in urine¹² and symmetric dimethylarginine in blood⁹) may follow to thoroughly evaluate an animal's renal condition.

Results of the present study indicated that straw-colored urine was most common and considered clinically normal for the Antillean manatees in the present study. The color is attributed to the presence of urochromes, uroerythrin, and urobilin that are formed through bilirubin metabolism and eliminated in the urine.²⁷ This urine color was consistent with that reported in other herbivorous species, such as cattle, goats, sheep, rabbits, rhinoceros (*Rhinoceros unicornis*, *Dicerorhinus sumatrensis*, and *Diceros bicornis*), and Floridian and Amazonian manatees^{8,26,28,29,b} but differed from the ochre or white colors of urine produced by horses and attributed to the presence of mucus and abundant amounts of squamous epithelial cells.³⁰ The straw-colored urine of manatees in the present report also differed from the typical yellow to dark-yellow urine of carnivores (including dogs, cats, dolphins, and pinnipeds), reflecting highly concentrated urine and high-protein diets.^{19,29,30} In the present study, the small proportion (4/32 [13%]) of manatees with urine that was colorless and had no alteration of other urinary pa-

rameters was similar to findings in Amazonian manatees in which dilute urine is thought to be associated with intake of large quantities of water.^{8,31} The urine of Antillean manatees sampled in the present study was predominantly clear, similar to urine from most domestic and wild mammals^{19,29,32,33} but unlike turbid urine of horses (owing to the presence of calcium carbonate crystals and mucus³⁰) and Amazonian manatees (owing to the presence of cells, crystals, and bacteria^{29,34}).

Urine odor varies among animals,²⁶ and a urinoid odor was most common in the urine of the Antillean manatees in the present study, consistent with findings for most mammals owing to the presence of volatile acids and decomposition products of urea.²⁷ Some manatees in the present study had urine with a grassy odor that was directly related to their herbivorous diet, and a few manatees from the Dominican Republic and Puerto Rico had urine that smelled fruity sweet, which we presumptively related to their dietary supplementation with fruits and vitamins. Although a fruity-sweet odor in urine could suggest glucosuria, none of the manatees sampled had glucose detected in their urine. Manatees under human care are fed diets rich in fats and carbohydrates,³⁵ and some animals have high serum fructosamine concentrations and possible hyperglycemia with type 2 diabetes.[†] In addition, previous studies^{10,b} indicate that urine glucose concentrations of 3.9 to 6.9 mg/dL and 3.0 to 3.6 mg/dL in Floridian and Amazonian manatees, respectively, possibly reflect the intake of carbohydrate-rich foods, such as beets.³⁰ Clinicians should pay close attention to urine glucose concentration in manatees.

Urinalysis results with test strips were negative for the detection of glucose, bilirubin, urobilinogen, ketones, nitrite, protein, blood, and leukocytes in the Antillean manatees of the present report. These findings were consistent with those in domestic animals,^{29,30} Asian elephants (*Elephas maximus*),⁸ rhinoceros,²⁸ common bottlenose dolphins (*Tursiops truncatus*),¹⁹ and Amazonian manatees.^{8,10} Similarly, the Antillean manatees in the present study did not have substantial numbers of RBCs or WBCs in their urine sediment when examined microscopically, as is typical of most domestic animals.³⁶ A previous study⁸ involving multiple urines samples from 21 Amazonian manatees with reactive urinalysis strips indicated most urine samples had 10 RBCs/mL and < 25 WBCs/mL; however, the investigators did not speculate on underlying causes, which could have included estrus in females or a urinary tract inflammation.³⁷ In the present study, microscopic evaluation of urine sediment revealed that the manatees had 0 to 8 WBCs/hpf and 0 to 3 RBCs/hpf, which were findings considered clinically normal because the cells are physiologically eliminated by the presentation of mild inflammation, heat, and epithelial desquamation.^{38,39} In contrast, Amazonian manatees have been reported⁸ to have various numbers of WBCs and RBCs and even suppurative exudate in their urine, suggesting the presence of genitourinary infections.

Marine mammals, such as common bottlenose dolphins, gray seals (*Halichoerus grypus*), and South American sea lions (*Otaria byronia*), tend to have acidic urine,^{8,19} and a previous study⁵ shows that Florida manatees also have acidic urine. However, our findings indicated that Antillean manatees have alkaline urine, and we hypothesized that this is a consequence of herbivory. Similarly, urine pH is neutral to alkaline (pH, 7.0 to 9.0) in other herbivorous mammals (eg, cattle, sheep, goats, horses, rabbits, rhinoceros, and elephants^{28,29,40}), likely owing to ingestion of large amounts of calcium carbonate, which is eliminated by renal excretion.²⁵

Our finding that the mean \pm SD urine SG was 1.010 ± 0.007 (range, 1.002 to 1.030) was consistent with reports^{41,42} that manatees have poorly concentrated urine under normal conditions as part of their physiologic osmoregulation process through semireniculated kidneys. Our findings suggested that the capacity to concentrate urine is similar between Antillean manatees and Amazonian manatees (urine SG, 1.000 to 1.015),⁸ Florida manatees (urine SG, 1.007 to 1.008),^{12,43,b} various domestic species, and herbivorous great apes^{44,45}; however, domestic and wild carnivores, cetaceans, and artiodactyls typically have more concentrated urine.^{19,29,37,46}

Urinary creatinine concentrations in the Antillean manatees of the present study were similar to those found in Amazonian and Florida manatees^{10,b}; cattle, sheep, and goats^{47,48}; domestic dogs⁴⁹; common bottlenose dolphins¹⁹; and harbor seals.⁵⁰ In addition, our findings for urinary microalbumin were also within reference limits for domestic mammals.^{37,49} Further, our findings for the mean microalbumin-to-creatinine ratio (100 mg/g) as an evaluation for early detection of renal dysfunction^{30,36,51} were similar to results in healthy dogs, cats, and other farm mammals,^{20,37,49,52,53} which suggested that the concentrations we observed were the result of physiologic excretion and clinically normal.

Epithelial cells were commonly observed in urinary samples from the Antillean manatees in the present study. This finding is clinically normal in mid-stream free-catch urine samples, including in Amazonian manatees,⁸ and occurs physiologically as a result of cell turnover.³⁴ Additionally, the finding of higher numbers of squamous epithelial cells in urine samples from females (vs males) may have been attributable to estrus cycles and associated accelerated detachment of vaginal epithelial cells that allowed a large number of these cells to be seen in the sediment of free-catch samples.^{8,54} The small numbers of transitional epithelial cells observed seemed to lack diagnostic importance, given that there were no associated abnormalities in other urine parameters.³⁸ Similarly, rare identification of RBC casts (only in 2 adult males) was considered a normal physiologic finding similar to that in domestic animals.^{34,37} In addition, our findings for urinary crystals were consistent with those in herbivores with alkaline urine (eg, horses,²⁹ Asian elephants,⁴⁰ rhinoceros,²⁸ and

Amazonian and Florida manatees^{8,55}) and other marine mammals (eg, common bottlenose dolphins,⁵⁶ ringed seals [*Pusa hispida*], northern elephant seals [*Mirounga angustirostris*], Weddell seals [*Leptonychotes weddellii*], and humpback whales [*Megaptera novaeangliae*]^{8,37}). The small amounts of bacteria and fungi or yeast observed in urine samples of the present study seemed to have represented contamination from free-catch sample collection in aquatic environments where complete sterility is difficult to obtain,^{37,38} particularly in male manatees owing to their long (up to approx 45 cm) tubular sheath within a prepuce that can harbor water from surroundings. In domestic and wild animals, the presence of fungal hyphae without accompanying abnormalities in other urine parameters is typically considered indicative of contamination, and such has been observed in Amazonian manatees.¹⁰ We also identified spermatozoa in the urine sample from 1 adult male and considered the finding clinically normal,^{30,34} as was our detection of mucus in urine from a male manatee.^{27,36} Although we did not stain slides for urinary sediment assessment because of logistic reasons owing to remote field circumstances in the present study, future studies would benefit from staining urine sediment slides and thereby making it easier to identify the sediment structures of interest.

Our finding of unidentified nematode-type larvae and eggs in the urine samples of 2 adult male Antillean manatees in Mexico was similar to that previously reported.⁵⁷ In contrast to mammalian herbivores, urinary sediment findings in carnivores and domestic omnivores more commonly include parasites (ie, giant dog kidney worm [*Diectophyma renale*], dog heartworm [*Dirofilaria immitis*], bladder worm [*Capillaria plica* and *Capillaria feliscati*], and swine kidney worm [*Sthepanurus dendatus*]^{33,37}). We suggest that future studies of Antillean manatees should isolate larvae and eggs found in urine samples and investigate their morphological and molecular properties to identify the organisms and ascertain whether they have any clinical impact on manatees.

Although the small number of manatees in each group when considered on the basis of age, sex, and habitat precluded further statistical analyses of results, to our knowledge, our overall findings yielded the first compilation of baseline values for urinalysis in healthy Antillean manatees under human care. We believe that the ease of free-catch urine collection methods (with operant conditioning and by opportunistic collection during physical examination) is encouraging for potential future expanded use in zoos, aquaria, or rescue centers for routine urinalysis in manatees and other wildlife species similarly considered difficult to sample for routine urinalysis. In addition, assessing symmetric dimethylarginine concentration in blood samples from Antillean manatees may serve as a relatively easy, fast, noninvasive, and cost-effective diagnostic tool to augment detection of renal disease or infection^a; additional research is warranted.

Acknowledgments

No third-party funding or support was received in connection with the present study or the writing or publication of the manuscript. The authors declare that there was no conflict of interest.

The authors thank A. M. Aguirre, T. Alejandro, L. Alvarado, M. R. Arreola, L. Bustamante, L. Garduño, N. Gutiérrez, N. Jiménez, A. Mingramm, F. Molina, Y. Moná, C. I. Rivera, S. Rivera, A. I. Rodríguez, and C. Rodríguez for assistance with animal training, handling, and urine collection; B. Rivera-Marchand for statistical analysis; and G. D. Bossart, K. Arevalo, J. C. Carrascal, M. Dennis, J. Gaspard, G. Melendez, and M. Walsh for assistance with the manuscript.

Footnotes

- a. Self-Sullivan C, Mignucci-Giannoni AA. *Trichechus manatus*. Gland, Switzerland: International Union for Conservation of Nature and Natural Resources [database online]. Available at: www.iucnredlist.org/species/22103/93569172008. Accessed January 16, 2021.
- b. Manire CA, Renner MS, Reidarson TH. Urinalysis and urine chemistries in the Florida manatee, *Trichechus manatus latirostris*, under differing diet and salinity (abstr), in *Proceedings*. 35th Annu Conf Int Assoc Aquatic Animal Med 2004.
- c. WL0020-ATC, Tekcoplus, Hong Kong.
- d. Mazuri Leaf-Eater Primate Diet, Land O'Lakes Inc, Arden Hills, Minn.
- e. Melesio-Navarro KM. *Técnicas para la realización de examen física general; pruebas de laboratorio y de gabinete en manatíes del Caribe (Trichechus manatus manatus) albergados en el Acuario de Veracruz A.C. mediante condicionamiento operante*. MVZ thesis, Universidad Veracruzana, Veracruz, Mexico, 2014.
- f. Urine collection container, Globe Scientific Inc, Mahwah, NJ.
- g. Mignucci-Giannoni AA, Jiménez-Zucchet AN, Cabrias-Contreras IJ. Marine mammal urinalysis. Centro de Conservación de Manatíes, 2020. Available at manatipr.org/wp-content/uploads/2020/07/MMUrinalysisForm.jpg. Accessed Jul 13, 2020.
- h. Excel, version 16.16.23, Microsoft Corp, Redmond, Wash.
- i. Nalgene (50 mL), Thermo Fisher Scientific, Waltham, Mass.
- j. Greiner Bio-One North America Inc, Monroe, NC.
- k. AimStrip 10-SG, Germain Laboratories, San Antonio, Tex.
- l. pH-Aware, HealthyWiser LLC, Wyo.
- m. Rapid Response Microalbumin 2-1 Combo Strip, BTNX Inc, Markham, ON, Canada.
- n. RETK-71, Tekcoplus Ltd, Hong Kong, China.
- o. Clay Adams Compact II centrifuge, Becton, Dickinson and Co, Franklin Lakes, NJ.
- p. SAS, version 9.1, SAS Institute Inc, Cary, NC.
- q. Smith LN, Bonde RK, Nau MR, et al. Significance of symmetric dimethylarginine in evaluating renal insufficiency in rehabilitated wild Florida manatees (*Trichechus manatus latirostris*) and reference values in two wild manatee populations (abstr), in *Proceedings*. 49th Annu Conf Am Assoc Zoo Vet 2017;68–69.
- r. Ball RL, Smith LN. Reference intervals of fructosamine in wild Florida manatees (*Trichechus manatus latirostris*) (abstr), in *Proceedings*. 50th Annu Conf Int Assoc Aquat Anim Med 2019.
- s. Kingsukon D, Soontornsit J, Prayurawongsa N, et al. Urinalysis in Asian elephants at Kochasuksa elephant study center, Thailand (abstr), in *Proceedings*. 2nd Symp Asian Zoo Wildl Med 1st Workshop Zoo Wildl Pathol (AZWMP) 2006;26–29.

References

1. Self-Sullivan C, Mignucci-Giannoni AA. West Indian manatees (*Trichechus manatus*) in the Wider Caribbean Region. In: Hines E, Reynolds J, Aragones L, et al, eds. *Sirenian conservation: issues and strategies in developing countries*. Gainesville, Fla: University Press of Florida, 2012;36–46.
2. Mignucci-Giannoni AA. Marine mammal captivity in the

- northeastern Caribbean, with notes on the rehabilitation of stranded whales, dolphins and manatees. *Caribb J Sci* 1998;34:191-203.
3. Bossart GD. Manatees. In: Dierauf LA, Gulland FM, eds. *CRC handbook of marine mammal medicine*. 2nd ed. Boca Raton, Fla: CRC Press, 2001;939-960.
 4. Bossart GD, Dierauf LA. Marine mammal clinical laboratory medicine. In: Dierauf LA, ed. *CRC handbook of marine mammal medicine: Health, disease, and rehabilitation*. Boca Raton, Fla: CRC Press, 1990;1-52.
 5. Bossart GD, Reidarson TH, Dierauf LA, et al. Clinical pathology. In: Dierauf LA, Gulland FM, eds. *CRC handbook of marine mammal medicine*. 2nd ed. Boca Raton, Fla: CRC Press, 2001;383-448.
 6. Davis MR, Walsh MT. Sirenian medicine. In: Gulland FMD, Dierauf LA, Whitman KL, eds. *CRC handbook of marine mammal medicine*. 3rd ed. Boca Raton, Fla: CRC Press, 2018;949-967.
 7. White JR, Francis-Floyd R. Manatee biology and medicine. In: Dierauf LA, ed. *CRC handbook of marine mammal medicine: health, disease, and rehabilitation*. Boca Raton, Fla: CRC Press, 1990;601-623.
 8. Pantoja TMdA, Rosas FCW, Da Silva VMF, et al. Amazonian manatee urinalysis: conservation applications. In: Povilitis T, ed. *Topics in conservation biology*. London: InTechOpen, 2012;57-80. Available at: www.intechopen.com/books/topics-in-conservation-biology. Accessed Jul 13, 2020.
 9. Manire CA, Walsh CJ, Rhinehart HL, et al. Alterations in blood and urine parameters in two Florida manatees (*Trichechus manatus latirostris*) from simulated conditions of release following rehabilitation. *Zoo Biol* 2003;22:103-120.
 10. Pantoja TMA, Rosas FCW, Da Silva VMF, et al. Urinary parameters of *Trichechus inunguis* (Mammalia, Sirenia): reference values for the Amazonian manatee. *Braz J Biol* 2010;70:607-615.
 11. Domning DP, Hayek LC. Interspecific and intraspecific morphological variation in manatees (Sirenia: *Trichechus*). *Mar Mamm Sci* 1986;2:87-144.
 12. Ortiz RM, Worthy GAJ, MacKenzie DS. Osmoregulation in wild and captive West Indian manatees (*Trichechus manatus*). *Physiol Zool* 1998;71:449-457.
 13. Vianna JA, Bonde RK, Caballero S, et al. Phylogeography, phylogeny and hybridization in trichechid sirenians: implications for manatee conservation. *Mol Ecol* 2006;15:433-447.
 14. Hunter ME, Mignucci-Giannoni AA, Pause Tucker K, et al. Puerto Rico and Florida manatees represent genetically distinct groups. *Conserv Genet* 2012;13:1623-1635.
 15. Mignucci-Giannoni AA, Montoya-Ospina RA, Jiménez-Marrero NM, et al. Manatee mortality in Puerto Rico. *Environ Manage* 2000;25:189-198.
 16. Colbert DE, Bauer GB. Basic husbandry training of two Florida manatees, *Trichechus manatus latirostris*. *Soundings* 1999;24:18-21.
 17. Colbert DE, Fellner W, Bauer GB, et al. Husbandry and research training of two Florida manatees (*Trichechus manatus latirostris*). *Aquat Mamm* 2001;27:16-23.
 18. Ramirez K. Marine mammal training: the history of training animals for medical behaviors and keys to their success. *Vet Clin North Am Exot Anim Pract* 2012;15:413-423.
 19. Jiménez-Zucchet N, Alejandro-Zayas T, Alvarado-Macedo CA, et al. Baseline urinalysis values in common bottlenose dolphins under human care in the Caribbean. *J Vet Diagn Invest* 2019;31:426-433.
 20. Sink CA, Weinstein NM. *Practical veterinary urinalysis*. Ames, Iowa: Wiley-Blackwell, 2012.
 21. Birukawa N, Ando H, Goto M, et al. Plasma and urine levels of electrolytes, urea and steroid hormones involved in osmoregulation of cetaceans. *Zoolog Sci* 2005;22:1245-1257.
 22. Daniels RL, Smith CR, Venn-Watson S. Effects of freeze-thaw cycle on urine values from bottlenose dolphins (*Tursiops truncatus*). *Aquat Mamm* 2013;39:330-334.
 23. Friedrichs KR, Harr KE, Freeman KP, et al. ASVCP reference interval guidelines: determination of de novo reference intervals in veterinary species and other related topics. *Vet Clin Pathol* 2012;41:441-453.
 24. Gunn-Christie RG, Flatland B, Friedrichs KR, et al. ASVCP quality assurance guidelines: control of preanalytical, analytical, and postanalytical factors for urinalysis, cytology, and clinical chemistry in veterinary laboratories. *Vet Clin Pathol* 2012;41:18-26.
 25. Rizzi TE, Vlenciano A, Bowles M, et al. *Atlas of canine and feline urinalysis*. Hoboken, NJ: John Wiley & Sons, 2017.
 26. Finco DR. Kidney function. In: Kaneko J, Harvey J, Bruss M, eds. *Clinical biochemistry of domestic animals*. Burlington, Mass: Academic Press, 2008;441-484.
 27. Campuzano-Maya G, Arbalaez-Gómez M. Uroanalysis: más que un examen de rutina. *Med Lab (Ed Ital)* 2006;12:511-555.
 28. Haffey MB, Pairan RD, Reinhart PR, et al. Urinalysis in three species of captive rhinoceros (*Rhinoceros unicornis*, *Dicerorhinus sumatrensis*, and *Diceros bicornis*). *J Zoo Wildl Med* 2008;39:349-357.
 29. Parrah JD, Moulvi BA, Gazi MA, et al. Importance of urinalysis in veterinary practice—a review. *Vet World* 2013;6:640-646.
 30. Bouda J, Doubek J, Quiroz Rocha G. Patología clínica del aparato urinario. In: Núñez-Ochoa L, Bouda J, eds. *Patología clínica veterinaria*. Mexico City: Universidad Nacional Autónoma, 2007;99-119.
 31. Díaz Portillo J, Fernández del Barrio MT, Paredes Salido F. *Aspectos básicos de bioquímica clínica*. Madrid, Spain: Ediciones Díaz de Santos, 1997.
 32. Barsanti JA, Finco DR. Laboratory findings in urinary tract infections. *Vet Clin North Am Small Anim Pract* 1980;9:729-748.
 33. Tripathi NK, Gregory CR, Latimer KS. Urinary system. In: Latimer KS, ed. *Duncan and Prasse's veterinary laboratory medicine: clinical pathology*. 5th ed. Ames, Iowa: Wiley-Blackwell, 2011;253-283.
 34. Callens AJ, Bartges JW. Urinalysis. *Vet Clin North Am Small Anim Pract* 2015;45:621-637.
 35. Siegal-Willott JL, Harr K, Hayek LC, et al. Proximate nutrient analyses of four species of submerged aquatic vegetation consumed by Florida manatee (*Trichechus manatus latirostris*) compared to romaine lettuce (*Lactuca sativa* var. *longifolia*). *J Zoo Wildl Med* 2010;41:594-602.
 36. King-Strasinger S, Schaub-Di-Lorenzo M. *Urinalysis and body fluids*. Philadelphia: FA Davis Co, 2014.
 37. Elliott J, Grauer GF, Westropp JL. *BSAVA manual of canine and feline nephrology and urology*. 3rd ed. Gloucester, England: British Small Animal Veterinary Association, 2017.
 38. Gómez-Lagos R, Pellegrini-Pinto P. *Recomendaciones para el análisis del sedimento urinario*. Santiago, Chile: Instituto de Salud Pública de Chile, 2013.
 39. Ruiz-Rodríguez J. Aproximación al análisis e bioquímica sanguínea y uroanálisis en animales silvestres y especies no convencionales. *Mem Conf Int Med Aprovech Fauna Silv Exót Conv* 2013;9:58-67.
 40. Wiedner E, Alleman AR, Isaza R. Urinalysis in Asian elephants (*Elephas maximus*). *J Zoo Wildl Med* 2009;40:659-666.
 41. Maluf NS. Renal anatomy of the manatee, *Trichechus manatus*, Linnaeus. *Am J Anat* 1989;184:269-286.
 42. Ortiz RM. Osmoregulation in marine mammals. *J Exp Biol* 2001;204:1831-1844.
 43. Irvine AB, Neal RC, Cardeilhac RT, et al. Clinical observations on captive and free-ranging West Indian manatees, *Trichechus manatus*. *Aquat Mamm* 1980;8:2-10.
 44. Sleeman JM, Mudakikwa AB. Analysis of urine from free-ranging mountain gorillas (*Gorilla gorilla beringei*) for normal physiologic values. *J Zoo Wildl Med* 1998;29:432-434.
 45. Kelly TR, Sleeman JM, Wrangham RW. Urinalysis in free-living chimpanzees (*Pan troglodytes schweinfurthii*) in Uganda. *Vet Rec* 2004;154:729-730.
 46. DelGiudice GD, Krausman PR, Bellantoni ES, et al. Blood and urinary profiles of free-ranging desert mule deer in Arizona. *J Wildl Dis* 1990;26:83-89.
 47. Bristow AW, Whitehead DC, Cockburn JE. Nitrogenous constituents in the urine of cattle, sheep and goats. *J Sci Food Agric* 1992;59:387-394.
 48. Chizzotti ML, Valaderes Filho SC, Valaderes RFD, et al. Determination of creatinine excretion and evaluation of spot urine sampling in Holstein cattle. *Livest Sci* 2008;113:218-225.

49. Cortadellas O, Fernández del Palacio MJ. Diagnóstico y tratamiento de la enfermedad renal crónica (ERC) en el perro y el gato. Parte 1: evaluación del paciente con ERC. *Clin Vet Pequeños Anim* 2012;32:215-223.
50. Smith HW. The composition of urine in the seal. *J Cell Comp Physiol* 1936;7:465-474.
51. Houston DM, Moore AEP. Canine and feline urolithiasis: examination of over 50,000 urolith submissions to the Canadian Veterinary Urolith Centre from 1998 to 2008. *Can Vet J* 2009;50:1263-1268.
52. Pressler BM, Vaden SL, Jensen WA, et al. Detection of canine microalbuminuria using semiquantitative test strips designed for use with human urine. *Vet Clin Patbol* 2002;31:56-60.
53. Whittemore JC, Gill VL, Jensen WA, et al. Evaluation of the association between microalbuminuria and the urine albumin-creatinine ratio and systemic disease in dogs. *J Am Vet Med Assoc* 2006;229:958-963.
54. Webb J. Renal cytology and urinalysis. In: Barger A, Macneill A, eds. *Small animal cytology diagnosis*. Boca Raton, Fla: CRC Press, 2016;357-378.
55. Keller M, Moliner JL, Vázquez G, et al. Nephrolithiasis and pyelonephritis in two West Indian manatees (*Trichechus manatus* spp.). *J Wildl Dis* 2008;44:707-711.
56. Sweeney JC, Reddy ML. Cetacean cytology. In: Dierauf LA, Gulland FM, eds. *CRC handbook of marine mammal medicine*. 2nd ed. Boca Raton, Fla: CRC Press, 2001;437-448.
57. Pires JML, Moreira AB, Rocha A. Pesquisa parasitária em urina de peixes-bois marinhos (*Trichechus manatus manatus*) mantidos em cativeiro no Brasil. *Sci Elec Arch* 2016;9:47-52.